

**RESEARCH ARTICLE**

# Clinical presentation and assessment of older patients presenting with headache to emergency departments: A multicentre observational study

Sierra Beck<sup>1,2</sup> | Frances B. Kinnear<sup>3,4</sup> | Anne Maree Kelly<sup>5,6</sup> |  
 Kevin H Chu<sup>7,8</sup> | Win Sen Kuan<sup>9,10</sup> | Gerben Keijzers<sup>11,12,13</sup> | Richard Body<sup>14,15</sup> |  
 Mehmet A. Karamercan<sup>16</sup> | Sharon Klim<sup>5</sup> | Tissa Wijeratne<sup>17,18,19</sup> |  
 Sinan Kamona<sup>20,21</sup> | Colin A Graham<sup>22</sup> | Tom Roberts<sup>23</sup> | Daniel Horner<sup>24,25</sup> |  
 Said Laribi<sup>26,27</sup> | HEAD Study Group

<sup>1</sup>Department of Medicine, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

<sup>2</sup>Emergency Department, Dunedin Hospital, Dunedin, New Zealand

<sup>3</sup>Emergency & Children's Services, The Prince Charles Hospital, Chermside, Qld, Australia

<sup>4</sup>Department of Medicine, University of Queensland, St Lucia, Qld, Australia

<sup>5</sup>Joseph Epstein Centre for Emergency Medicine Research @ Western Health, Sunshine, Melbourne, Vic., Australia

<sup>6</sup>Department of Medicine - Western Health, Melbourne Medical School, The University of Melbourne, Parkville, Vic., Australia

<sup>7</sup>Emergency and Trauma Centre, Royal Brisbane and Women's Hospital, Herston, Qld, Australia

<sup>8</sup>Faculty of Medicine, University of Queensland, St Lucia, Qld, Australia

<sup>9</sup>Emergency Medicine Department, National University Hospital, National University Health System, Singapore

<sup>10</sup>Department of Surgery, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

<sup>11</sup>Department of Emergency Medicine, Gold Coast University Hospital, Gold Coast, Qld, Australia

<sup>12</sup>Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Qld, Australia

<sup>13</sup>School of Medicine, Griffith University, Gold Coast, Qld, Australia

<sup>14</sup>Division of Cardiovascular Sciences, The University of Manchester, Manchester, UK

<sup>15</sup>Emergency Department, Manchester Academic Health Science Centre, Manchester University NHS Foundation Trust, Manchester, UK

<sup>16</sup>Faculty of Medicine Department of Emergency Medicine, Gazi University, Ankara, Turkey

<sup>17</sup>The University of Melbourne, Parkville, Australia

<sup>18</sup>Department of Neurology, Western Health, St Albans, Vic., Australia

<sup>19</sup>Public Health School, La Trobe University, Bundoora, Vic., Australia

<sup>20</sup>School of Medicine, University of Auckland, Auckland, New Zealand

<sup>21</sup>Auckland District Health Board, Auckland City Hospital, Auckland, New Zealand

<sup>22</sup>Emergency Medicine, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, New Territories, Hong Kong SAR

<sup>23</sup>Trainee Emergency Research Network (TERN), North Bristol NHS Trust, Bristol, UK

<sup>24</sup>Emergency Department, Salford Royal NHS Foundation Trust, Stott Lane, Salford, UK

<sup>25</sup>Division of Infection, Immunity and Respiratory Medicine, University of Manchester, Oxford, UK

<sup>26</sup>Emergency Medicine Department, Tours University Hospital, Tours, France

<sup>27</sup>EUSEM Research Network, Aartselaar, Belgium

**Clinical trials registration:** The study was registered with the Australia and New Zealand Clinical Trials Register (trial number 376 695)

**Correspondence**

Anne Maree Kelly, Joseph Epstein Centre for Emergency Medicine Research @ Western Health, Sunshine, Melbourne, Vic., Australia.  
Email: Anne-Maree.Kelly@wh.org.au

**Funding information**

Royal College of Emergency Medicine (UK) provided part funding.

**Abstract**

**Objective:** To describe the characteristics, assessment and management of older emergency department (ED) patients with non-traumatic headache.

**Methods:** Planned sub-study of a prospective, multicentre, international, observational study, which included adult patients presenting to ED with non-traumatic headache. Patients aged  $\geq 75$  years were compared to those aged  $< 75$  years. Outcomes of interest were epidemiology, investigations, serious headache diagnosis and outcome.

**Results:** A total of 298 patients (7%) in the parent study were aged  $\geq 75$  years. Older patients were less likely to report severe headache pain or subjective fever (both  $P < 0.001$ ). On examination, older patients were more likely to be confused, have lower Glasgow Coma Scores and to have new neurological deficits (all  $P < 0.001$ ). Serious secondary headache disorder (composite of headache due to subarachnoid haemorrhage (SAH), intracranial haemorrhage, meningitis, encephalitis, cerebral abscess, neoplasm, hydrocephalus, vascular dissection, stroke, hypertensive crisis, temporal arteritis, idiopathic intracranial hypertension or ventriculoperitoneal shunt complications) was diagnosed in 18% of older patients compared to 6% of younger patients ( $P < 0.001$ ). Computed tomography brain imaging was performed in 66% of patients  $\geq 75$  years compared to 35% of younger patients ( $P < 0.001$ ). Older patients were less likely to be discharged (43% vs 63%,  $P < 0.001$ ).

**Conclusion:** Older patients with headache had different clinical features to the younger cohort and were more likely to have a serious secondary cause of headache than younger adults. There should be a low threshold for investigation in older patients attending ED with non-traumatic headache.

**KEYWORDS**

aged, emergency departments, epidemiology, headache

## 1 | INTRODUCTION

Headache makes up a significant proportion of emergency department (ED) visits.<sup>1-3</sup> In a recent population-based United States (US) study, headache was the fifth most common reason to present to EDs, representing 3% of all visits.<sup>1</sup> Little is known about the epidemiology of older patients who present to the ED with headache. Prior neurology-based studies have shown the incidence of patients presenting with headache decreases with age, but the risk of headache due to serious conditions increases.<sup>4-6</sup> The prevalence of migraine is known to decrease with age<sup>4-7</sup> and likely contributes to this decline in total headache presentations. Concern that a headache may represent a secondary headache disorder caused by a serious condition such as stroke, temporal arteritis or intracranial malignancy, increases in older populations. In a previous

**PRACTICE IMPACT**

In this international multicentre study, a serious headache disorder was diagnosed in 18% of patients aged  $\geq 75$  years compared to 6% of younger patients. Older patients had higher imaging utilisation rates, receiving a computed tomography scan of the brain in 66% of presentations compared to 35% of younger patients. In contrast to younger patients, patients aged  $\geq 75$  years were more likely to report mild pain, and less frequently reported associated fevers, nausea, vomiting and photophobia. There should be a low threshold for investigation in older patients attending ED with non-traumatic headache.

single-centre study of patients aged >65 years referred to a neurology service with new onset headache, 15% had a secondary headache disorder with a serious underlying cause.<sup>6</sup>

Increasing age is cited as a 'red flag' for the identification of serious secondary headache disorders in the ED.<sup>8-12</sup> Emergency medicine research supports age over 40 as a risk factor for subarachnoid haemorrhage.<sup>9</sup> Age over 50 has been identified as a risk factor for the identification of serious secondary causes of headache.<sup>10-13</sup> Reflecting this evidence, clinical guidelines recommend that clinicians consider diagnostic work-up of headaches for severe underlying cause in patients with increasing age.<sup>8,14,15</sup> Little is known regarding the ways that older people's headache signs and symptoms may differ from younger patients. Anticoagulation use, polypharmacy and increased comorbidities further complicate the diagnostic work-up and management of older patients. There is a shortage of real-world data specific to ED patients describing the differences in clinical characteristics, management and outcomes of older patients with headache.

Generally, the objective of the Headache in Emergency Departments (HEAD) study was to describe the epidemiology of non-traumatic headache in adults presenting to the ED including clinical presentation, investigations and disposition.<sup>16</sup> In this planned sub-study, we focus on patients aged 75 years or older and their comparison with patients <75 years with headache. This group is under-represented in research and is likely to have specific characteristics, which may impact their care. We hypothesised that this study would confirm previous research that serious secondary causes of headache are more common in older people.

## 2 | METHODS

### 2.1 | Design and settings

This was a planned sub-study of a prospective multi-centre observational study conducted over one calendar month in 2019, the Headache in Emergency Departments (HEAD) Study.<sup>16</sup>

### 2.2 | Participants and recruitment

For the parent study, participants were adult patients (aged ≥18 years) with non-traumatic headache as their presenting complaint. Exclusion criteria were history of trauma to the head within 48 hours of presentation, missing records, inter-hospital transfers, re-presentation with the same headache as at a recent previous visit and headache

as an associated symptom rather than a main complaint. Determination of whether headache was a primary complaint was at the discretion of the local researcher based on all available data.

Qualifying adult patients presenting during the study period were identified from ED data management systems. Patients were identified prospectively but, depending on site resources, some data were collected retrospectively.

### 2.3 | Data collected and data collection process

Data collected included demographics, clinical assessment, investigation, treatment, diagnosis, disposition and outcome. Data were collected from clinical records by local researchers onto piloted data forms or directly into the study database (depending on local processes and resources). Study data were collected and managed using REDCap electronic data capture tools hosted at the Joseph Epstein Centre for Emergency Medicine Research [Melbourne, Australia]. For the majority of sites, the data were collected in March 2019, with some minor variations following delays in ethical/governance approval.

### 2.4 | Outcomes of interest

The primary outcomes of interest of this sub-study were the differences in epidemiology, investigations, serious secondary headache diagnosis and disposition of patients aged 75 years or older presenting to the ED with headache. We chose 75 years as the cut-off consistent with recent recommendations.<sup>17,18</sup>

Severe headache was defined as pain score 7-10. Thunderclap headache was defined as headache peaking instantly or almost instantly. This was distinguished from headache that reached peak intensity within one hour but not instantly.

We defined non-migraine benign headache as the composite of those classified as primary, tension, cluster or musculoskeletal following clinical assessment and initial management. Migraine is reported separately. As in the parent study, serious secondary headache disorders were defined as the composite of headache due to subarachnoid haemorrhage (SAH), intracranial haemorrhage, meningitis, encephalitis, cerebral abscess, neoplasm, hydrocephalus, vascular dissection, stroke, hypertensive crisis or pregnancy-related hypertension, temporal arteritis, idiopathic intracranial hypertension or ventriculoperitoneal shunt complications, noting that pregnancy-related hypertension would not apply to patients aged >75 years.

## 2.5 | Analysis and sample size

As this was a descriptive study, a formal sample size calculation was not performed. Descriptive analysis was performed using indicators of central tendency in normally distributed variables and frequencies in discrete variables. Non-parametric analyses were used as required. Comparison of proportions was analysed by chi square or Fisher's test as appropriate. The study is reported in accordance with STROBE guidelines for observational work.<sup>19</sup> Because this study was a sub-study, no a priori sample size was calculated. Post hoc power calculation was undertaken using University of British Columbia online power/sample size tool.<sup>20</sup> For the endpoint of serious secondary headache, to identify

a difference in rate of 7% (ie a doubling from 7% in the parent study<sup>16</sup> to 14%) with  $\alpha$  of 0.05 and power of 0.8 required 129 patients.

## 2.6 | Ethics approval

The lead ethics approval was by Melbourne Health Human Research Ethics Committee (HREC/43148/MH-2018). Ethics approval was subsequently obtained for each participating site according to local institutional requirements. In most jurisdictions, the study was conducted under waiver of consent. Patient consent was required in a few jurisdictions. In some Queensland sites, formal consent was required. This was verbal. In the United Kingdom

TABLE 1 Patient demographics

Variable	Result (Total N = 4536)	Age <75 years (4238, 93%)	Age ≥75 years (298, 7%)	Missing data (N)	P value
Age (years, median, IQR)	41 (29-55)	39 (28-52)	81 (77-86)	0	N/A
Gender (male, N, %)	1627, 36%	1518, 36%	109, 37%	1	0.002
Referral by a doctor (N, %)	788, 17%	722, 17%	66, 22%	0	0.02
Arrival by ambulance (N, %)	791, 17%	654, 15%	137, 46%	0	<0.001
Duration of symptoms (N, %)					
<24h	2060, 46%	1910, 46%	150, 51%	70	0.02 <sup>a</sup>
1-3 days	1021, 23%	975, 23%	46, 15%		
>3 days	1385, 31%	1289, 31%	96, 33%		
Triage category (N, %)					
Immediate	77, 2%	66, 2%	11, 4%	0	<0.001 <sup>a</sup>
Urgent	2294, 51%	2103, 50%	191, 64%		
Non-urgent	2165, 48%	2069, 49%	96, 32%		
Past medical history (N, %)					
Physician-diagnosed migraine	953, 21%	934, 22%	19, 6%	0	<0.001
Stroke/TIA	144, 3%	92, 2%	52, 17%	0	<0.001
Known cerebral malignancy-primary or secondary	44, 1%	43, 1%	1, 0.3%	0	0.3
Active non-cerebral malignancy	67, 1%	51, 1%	16, 5%	0	<0.001
Serious intracranial injury	48, 1%	41, 1%	7, 2%	0	0.1
Ventriculoperitoneal shunt	42, 1%	40, 1%	2, 1%	0	0.7
Intracranial aneurysm without haemorrhage	38, 1%	27, 1%	11, 4%	0	0.001
Subarachnoid haemorrhage	19, 0.4%	16, 0.4%	3, 1%	0	0.2
Known vascular abnormality	18, 0.4%	16, 0.4%	2, 1%	0	0.7
Chronic Medications (N, %)					
Opioids (excluding codeine)	115, 3%	104, 2%	11, 4%	0	0.1
Codeine preparations	64, 1%	56, 1%	8, 3%	0	0.7
Long acting anticoagulants	102, 2%	50, 1%	52, 17%	0	<0.001

Abbreviations: IQR, interquartile range; N, number; TIA, transient ischaemic attack.

<sup>a</sup>Omnibus Chi square.

(UK), an opt-out approach was used, and approval obtained through the Health Research Authorisation following application and review by committee (REC reference: 19/SW/0089).

The study was prospectively registered with the Australia and New Zealand Clinical Trials Register (trial number 376 695).

### 3 | RESULTS

A total of 4,536 patients presenting with headache were included in the parent study across 67 hospital groups (74 EDs) in 10 countries (Australia 34, New Zealand

10, Turkey 9, UK 7, Singapore 4, Belgium, 4, France 3, Romania 1, Hong Kong 1, Israel 1). For hospitals that provided caseload data, headache patients accounted for 1% of ED caseload (95% CI 0–1%). Two hundred and ninety-eight (7%) of the 4,536 patients were aged 75 years and over. The distribution by country is summarised in Appendix S1.

Patient demographics are summarised in Table 1. There was no difference in the proportion referred by a doctor or the duration of symptoms prior to presentation by age cohort. The older patient group was approximately three times more likely to arrive by ambulance. There were significant differences in triage category allocation with immediate or urgent triage categories more frequently

TABLE 2 Clinical symptoms

Clinical symptoms (N, %)	Total (N = 4536)	Age <75 years (N = 4238)	Age ≥75 years (N = 298)	Missing data (N)	P value
Onset of symptoms <sup>a</sup>					
Gradual	2486, 71%	2341, 71%	145, 67%	0	0.3 <sup>b</sup>
Sudden/thunderclap (instant peak)	644, 18%	595, 18%	49, 23%		
Peak intensity <1 hour	366, 10%	344, 10%	22, 10%		
Unknown	1040 (23%) <sup>a</sup>	958 (23%) <sup>a</sup>	82 (28%) <sup>a</sup>		
Location of headache <sup>a</sup>					
Generalised	2597, 67%	2429, 67%	168, 68%	0	0.4 <sup>b</sup>
Unilateral	1290, 33%	1210, 33%	80, 32%		
Unclear	648 (14%) <sup>a</sup>	598 (14%) <sup>a</sup>	50 (17%) <sup>a</sup>		
Severity <sup>a</sup>					
Mild (pain score ≤3)	815, 21%	742, 20%	73, 31%	0	<0.001 <sup>b</sup>
Moderate (pain score 4-7)	1869, 48%	1766, 48%	103, 44%		
Severe (pain score >7)	1235, 32%	1177, 32%	58, 25%		
Unclear	616 (14%) <sup>a</sup>	552 (13%) <sup>a</sup>	64 (21%) <sup>a</sup>		
Associated Symptoms					
Reported neck pain or stiffness	756, 17%	721, 17%	35, 12%	0	0.02
Nausea or vomiting	1844, 41%	1767, 42%	77, 26%	0	<0.001
Syncope/loss of consciousness	136, 3%	126, 3%	10, 3%	0	0.7
Reported photophobia	940, 21%	920, 22%	20, 7%	0	<0.001
New limb weakness (current/transient)	201, 4%	178, 4%	23, 8%	0	0.004
New limb paraesthesia (current/transient)	268, 6%	255, 6%	13, 4%	0	0.2
New speech difficulty (current/transient)	152, 3%	132, 3%	20, 7%	0	<0.001
New visual disturbance (current/transient)	600, 13%	560, 13%	40, 13%	0	0.9
Subjective fever or rigours	406, 9%	398, 9%	8, 3%	0	<0.001
Reported rash	56, 1%	50, 1%	6, 2%	0	0.3
Current or recent IV drug use	30, 1%	29, 1%	1, 0.3%	0	0.7
Worst headache ever	582, 13%	554, 13%	28, 9%	0	0.07
Head trauma within the last week	111, 2%	99, 2%	12, 4%	0	0.07

Abbreviation: IV, intravenous.

<sup>a</sup>For named categories reported as per cent of total with a named category response. For unknown/unclear reported as per cent of total.

<sup>b</sup>Omnibus Chi square.

TABLE 3 Clinical examination

Clinical examination	Total (N = 4536) (N, %)	Age <75 years (N = 4238) (N, %)	Age ≥75 years (N = 298) (N, %)	Missing data N, (N age <75 years, age ≥75 years)	P value
Vital signs					
Pulse rate >110 (N, %)	149, 3%	144, 3%	5, 2%	0	0.09
Systolic BP >160 mmHg	540, 12%	485, 12%	116, 9%	63, 1% (60, 1%/3, 1%)	<0.001
Systolic BP <90 mmHg	8, 0.2%	8, 0.2%	0		NS
Temperature >38C (N, %)	104, 3%	102, 3%	2, 1%	632, 14% (596, 14%/36, 12%)	0.07
Glasgow Coma Score (N, %)				537	
15	3921, 98%	3681, 98%	240, 93%	0	<0.001 <sup>a</sup>
13-14	62, 2%	46, 1%	16, 6%		
10-12	8, 0.2%	6, 0.2%	2, 1%		
<10	8, 0.2%	7, 0.2%	1, 0.4%		
Any new neurological signs on examination					
Isolated speech deficit	147, 3%	124, 3%	23, 8%	0	<0.001
Isolated motor deficit	10, 0.2%	8, 0.2%	2, 1%		
Speech and motor deficit	46, 1%	40, 1%	6, 2%		
Incoordination/ cerebellar signs	7, 0.2%	5, 0.1%	2, 1%		
Incoordination/ cerebellar signs	24, 0.5%	20, 0.5%	4, 1%		
New visual defect on examination	97, 2%	87, 2%	10, 2%		
Other exam findings					
Rash on examination	70, 2%	63, 2%	7, 2%	0	0.2
Confusion on examination	68, 1%	52, 1%	16, 5%	0	<0.001
Meningism on examination	63, 1%	60, 1%	3, 1%	0	0.8

Abbreviation: BP, blood pressure.

<sup>a</sup>Omnibus Chi square.

assigned to patients in the older age group. Older patients were also more likely to have a past history of stroke/transient ischaemic attack (TIA), active non-cerebral malignancy, intracranial aneurysm without haemorrhage and long-acting anticoagulant use. A past history of physician-diagnosed migraine was less common in the older age group (6% vs 22%).

Clinical symptoms are summarised in Table 2. Onset of symptoms and headache location proportions were similar between the age groups as were moderate pain scores and the percentage of participants who described their headache as their worst ever. Severe pain was less common, and mild pain was more common in the older group compared to younger patients. Nausea or vomiting, reported photophobia, and subjective fever or rigours were all also less common in older patients. Patient-reported new limb weakness or new speech difficulty were more common in the older age group. Syncope and reported neck pain or stiffness were distributed equally.

Examination findings are summarised in Table 3. New neurological signs (8% vs 3%), lower Glasgow Coma Score (GCS) and confusion (5% vs 1%) were more common in older patients. Demonstrable meningism was distributed equally.

Diagnostic imaging investigations, dispositions and outcomes are summarised in Table 4. Computed tomography (CT) of the brain was performed almost twice as often in the older age group (66% vs 35%). Lumbar puncture was performed much less frequently in older patients. There was no difference between the age groups with respect to magnetic resonance imaging (MRI), CT angiography, CT plus MRI or CT plus CT angiography, all of which were uncommonly performed (all less than 5% of total patients). More patients in the older age group required hospital admission.

The final ED diagnosis is shown in Table 5. With respect to specific diagnoses, migraine was less common in the older patient group compared to younger patients.



TABLE 4 Diagnostics imaging investigations, dispositions and outcomes

Investigations	Total (N = 4536)	Age <75 years (N = 4238)	Age ≥75 years (N = 298)	Missing data (N)	P value
	(N, %)	(N, %)	(N, %)		
Brain CT	1661, 37%	1471, 35%	190, 66%	0	<0.001
Normal	1360, 82%	1222, 83%	138, 73%		<0.001
SAH	31, 2%	27, 2%	3, 2%		NS
Other intracranial bleed	57, 3%	39, 3%	14, 7%		<0.001
Abscess	0	0	0		
Neoplasm	31, 2%	28, 2%	3, 2%		NS
MRI	151, 3%	138, 3%	13, 4%	0	0.3
Normal	89, 59%	84, 62%	5, 38%		0.1
Intracranial bleed	3, 2%	2, 1%	1, 8%		
Abscess	1, 1%	0	1, 8%		
Neoplasm	16, 11%	15, 11%	1, 8%		
Other (incl. non-acute changes)	42, 28%	37, 27%	5, 38%		
CT angiography	219, 5%	195, 5%	24, 8%	0	0.007
Normal	158, 72	141, 72%	17, 71%		
Aneurysm with bleed	19, 9%	18, 9%	1, 4%		0.3
Aneurysm without bleed	8, 4%	6, 3%	2, 8%		0.2
Lumbar puncture	171, 4%	169, 4%	2, 0.3%	0	0.001
CT plus MRI	114, 3%	104, 2%	10, 3%	0	0.9
CT plus CTA	196, 4%	175, 4%	21, 7%	0	0.02
<b>Disposition</b>	<b>(N, %)</b>	<b>(N, %)</b>	<b>(N, %)</b>	<b>0</b>	
Home from ED (including from ED observation unit)	3792, 84%	3606, 85%	186, 62%		<0.001 <sup>a</sup>
Ward	633, 14%	537, 13%	96, 32%		
Critical care	33, 1%	31, 1%	2, 1%		
Transfer	63, 1%	51, 1%	12, 4%		
Died in ED	1, 0.02%	0	1, 0.3%		
Operating theatre	3, 0.07%	3, 0.07%	0		
Interventional radiology	1, 0.02%	0	1, 0.3%		
Unknown	10, 0.2%	10, 0.2%	0		
<b>In hospital outcome</b>	<b>(N, %, CI)</b>	<b>(N, %, CI)</b>	<b>(N, %, CI)</b>	<b>0</b>	
Mortality	11, 0.2%	7, 0.2%	4, 1%		0.07

Abbreviations: CT, computed tomography; CTA, computed tomography angiogram; ED, emergency department; MRI, magnetic resonance imaging; SAH, subarachnoid haemorrhage.

<sup>a</sup>Omnibus Chi square.

Stroke/transient ischaemic attack and non-subarachnoid haemorrhage were proportionately more frequent in the 75 years and over age group. Serious secondary headache disorders, as defined above, were identified in 7% of patients overall, but were three times more common in older patients. In the ≥75 years age group, reported severity of pain was not associated with a serious secondary cause ( $P = 0.5$ ).

## 4 | DISCUSSION

In this study, in contrast to younger patients, patients aged ≥75 years were more likely to report mild pain, new limb weakness or speech difficulties, and less frequently reported associated fever, nausea, vomiting and photophobia. They were more likely to have a serious cause for their headache diagnosed in the ED. They also had higher

TABLE 5 Final ED diagnosis

ED diagnosis	Total (N = 4536)	Age <75 years (N = 4238)	Age ≥75 years (N = 298)	P value (major selected)
Non-migraine primary headache (unspecified)	1598, 35%	1491, 35%	107, 36%	0.8
Migraine	1101, 24%	1088, 26%	13, 4%	<0.001
Tension headache	317, 7%	296, 7%	21, 7%	
Viral illness (non-meningitis)	204, 4%	199, 5%	5, 2%	
Sinusitis	141, 3%	139, 3%	2, 1%	
Post-traumatic headache	76, 2%	70, 2%	6, 2%	
Musculoskeletal	72, 2%	66, 2%	6, 2%	
Cluster headache	71, 2%	71, 2%	0	
Stroke/ TIA	68, 1%	46, 1%	22, 7%	<0.001
Stroke <sup>a</sup>	50, 1%	33, 1%	17, 6%	
TIA	18, 0.4%	13, 0.3%	5, 2%	
Hypertension	64, 1%	51, 1%	13, 4%	<0.001
Hypertension crisis/urgency/malignant <sup>a</sup>	11, 0.2%	9, 0.2%	2, 1%	
Pregnancy-related hypertension <sup>a</sup>	5, 0.1%	5, 0.1%	0	
Hypertension-other	48, 1%	37, 1%	11, 4%	
Non-SAH intracranial haemorrhage / haematoma <sup>a</sup>	57, 1%	41, 1%	16, 5%	<0.001
Subarachnoid haemorrhage <sup>a</sup>	44, 1%	39, 1%	5, 2%	
Neoplasm <sup>a</sup>	43, 1%	38, 1%	5, 2%	
Meningitis (all)	48, 1%	48, 1%	0	
Viral <sup>a</sup>	40, 1%	40, 1%	0	
Bacteria <sup>a</sup>	3, 0.07%	3, 0.07%	0	
Unknown <sup>a</sup>	5, 0.1%	5, 0.1%	0	
Trigeminal neuralgia/cranial neuralgia	34, 1%	26, 1%	8, 3%	
Non-cranial sepsis (eg pneumonia, UTI, tonsillitis)	29, 1%	28, 1%	1, 0.3%	
Idiopathic Intracranial hypertension <sup>a</sup>	27, 1%	26, 1%	1, 0.3%	
Vertigo/BPPV	23, 0.5%	18, 0.4%	5, 2%	
Postlumbar puncture headache	13, 0.3%	13, 0.3%	0	
Ventriculoperitoneal shunt issues <sup>a</sup>	12, 0.3%	12, 0.3%	0	
Temporal arteritis <sup>a</sup>	11, 0.2%	6, 0.1%	5, 2%	
Dental cause	10, 0.2%	10, 0.2%	0	
Anxiety/psychogenic	9, 0.2%	8, 0.2%	1, 0.3%	
Alcohol-related hangover	8, 0.2%	8, 0.2%	0	
Postcoital headache	8, 0.2%	8, 0.2%	0	
Aneurysm/vascular malformation	8, 0.2%	8, 0.2%	0	
Toxicity	6, 0.1%	6, 0.1%	0	
Hyponatraemia	6, 0.1%	4, 0.09%	2, 1%	
Herpes zoster of head and neck	6, 0.1%	3, 0.07%	3, 1%	
Encephalitis <sup>a</sup>	6, 0.1%	5, 0.1%	1, 0.3%	
Seizure	6, 0.1%	6, 0.1%	0	
Vascular dissection <sup>a</sup>	4, 0.09%	4, 0.09%	0	
Hydrocephalus <sup>a</sup>	4, 0.09%	3, 0.07%	1, 0.3%	
Analgesia overuse syndrome	2, 0.04%	2, 0.05%	0	
Glaucoma	2, 0.04%	2, 0.05%	0	

(Continues)



TABLE 5 (Continues)

ED diagnosis	Total (N = 4536)	Age <75 years (N = 4238)	Age ≥75 years (N = 298)	P value (major selected)
Cerebral abscess <sup>a</sup>	1, 0.02%	1, 0.02%	0	
Other	157, 3%	134, 3%	23, 8%	
Unclear	240, 5%	214, 5%	26, 9%	
Serious secondary cause (marked <sup>a</sup> )	323, 7%	245, 6%	53, 18%	<0.001

Abbreviations: BPPV, benign paroxysmal positional vertigo; SAH, subarachnoid haemorrhage; TIA, transient ischaemic attack; UTI, urinary tract infection.

<sup>a</sup>Omnibus Chi square analysis.

imaging utilisation and admission rates. The prevalence of headaches associated with a serious underlying cause identified in this study (18%) is important because it supports and validates data from previous neurology literature.<sup>6,13</sup> Our results are also in keeping with prior literature in non-ED settings demonstrating lower incidence and prevalence of migraine with age.<sup>4,5,7</sup>

The high rate of CT scan in older patients with headache is noteworthy. While this will be in part due to higher prevalence of new neurological findings or confusion on examination, it may also reflect an awareness by clinicians that, as found in this study, about 1 in 5 older people presenting with headache will have a serious secondary cause. The high rate of serious secondary cause reinforces that clinicians should have a low threshold for investigation, especially neuroimaging.

As a group, older patients with headache presented with less severe pain and fewer associated general non-neurological symptoms. Severity of pain was not associated with identification of a serious secondary cause. Possible explanations include changes in disease prevalence with age (particularly the decreasing prevalence of migraine), immunosenescence, physiologic changes of ageing or underreporting and recording of symptoms.<sup>7,21,22</sup> Older adults are known to present atypically or with less severe symptoms for a variety of disease processes including myocardial infarction, urinary tract infection, intra-abdominal infection and traumatic intracranial haemorrhage.<sup>23-26</sup>

Previous findings in older patients with traumatic intracranial haemorrhage have parallels to our work on non-traumatic headache. Older patients presenting with traumatic intracranial haemorrhage were shown to have a higher level of consciousness (based on Glasgow Coma Scale score) compared to younger patients for a given anatomical severity of injury.<sup>26</sup> Some proposed reasons for this include increased intracranial capacity secondary to cerebral atrophy, differing progression of pathology and less obvious impact on higher cognitive function.<sup>26</sup> Certain clinical indicators of severe disease can be subtle and

harder to detect in older patients, making them potentially more vulnerable to missed or delayed diagnosis. Further research exploring whether older patients with serious secondary causes of headache exhibit different clinical features than a younger cohort may warrant exploration.

Our study has strengths in its size and geographical variation. Our study has some limitations, which should be considered when interpreting the data. While our study comprised a large number of patients with headache-related ED presentations from a range of countries, hospital sizes and medical systems, four of the ten countries accounted for approximately 85% of the presentations. This may in part reflect how communities use EDs. It is possible that some diagnoses are region dependent. In addition, access to and ordering of neuroimaging may vary between countries.<sup>16,27</sup> Data abstraction in this study was performed by local researchers at each site supported by a data dictionary but not formal training, which may have compromised the quality of the source data. Cross-checking of data abstraction was not performed. We did not collect data about inclusion/exclusion response, which may limit the generalisability of our findings. Data collection forms were however standardised and piloted. While attempts were made to prospectively enrol patients and collect data, this was not feasible at all sites because of resource constraints; retrospective data entry may have further limited data quality and increased the proportion of missing data.<sup>28</sup> The final ED diagnosis was recorded based on a subjective clinical impression derived from information available at the point of ED discharge. Additional information later may have revealed an alternative pathological diagnosis. We did not collect data on functional baseline, or goals of care status which may have impacted decisions regarding diagnostic work-up in older patients and subsequent clinical impression at discharge. Despite the overall large numbers in the study, the range of diagnoses represented in this cohort has resulted in low absolute numbers for certain diagnoses. As such, generalisability is limited.

## 5 | CONCLUSIONS

In this study, patients aged  $\geq 75$  years were more likely than younger patients to report mild pain, new limb weakness or speech difficulties and less frequently reported associated fever, nausea, vomiting and photophobia. On examination, they were more likely to be confused and have new neurological signs. Older patients with headache were more likely to have a serious secondary cause for their headache identified in the ED. Imaging and management pathways should reflect these increased risks.

### ACKNOWLEDGEMENTS

**The authors would like to thank the HEAD study group:** Catherine Lunter (Coffs Harbour Hospital, New South Wales, Australia), Rochelle Facer (Concord Repatriation Hospital, New South Wales, Australia), David Thomson (Port Macquarie Base and Kempsey District Hospitals, New South Wales, Australia), Robert Day (Royal North Shore Hospital, New South Wales, Australia), Greg McDonald (Sydney Adventist Hospital, New South Wales, Australia), Sarah Jones (Tamworth Regional Hospital, New South Wales, Australia), Julian Cochrane (Orange Base Hospital, New South Wales, Australia), Stephen Gourley (Alice Springs Hospital, Northern Territory, Australia), Mark Ross & Vinay Gangathimmaiah (Royal Darwin Hospital, Northern Territory, Australia), Kim Hansen (St Andrew's War Memorial Hospital, Queensland, Australia); Frances B Kinnear (The Prince Charles Hospital, Queensland, Australia), Gerben Keijzers (Gold Coast University Hospital, Queensland, Australia), Kevin Chu (Royal Brisbane and Women's Hospital, Queensland, Australia), Paul Bowe (Robina Hospital, Queensland, Australia), Raymund de la Cruz (Lyell McEwin and Modbury Hospitals, South Australia, Australia), Daniel Haustead (The Queen Elizabeth and Royal Adelaide Hospitals, South Australia, Australia), Jean Moller (University Hospital Geelong, Victoria, Australia), Katie Walker (Cabrini Malvern, Victoria, Australia), Richard D Smith (Bendigo Health, Victoria, Australia), Ron Sultana (Epworth Healthcare, Victoria, Australia), John Pasco (Werribee Mercy Hospital, Victoria, Australia), Neil Goldie and Andis Gaudins (Monash Health, Victoria, Australia), Rosamond Dwyer (Peninsula Health, Victoria, Australia), George Plunkett (Melbourne Health, Victoria, Australia); Anne-Maree Kelly (Western Health, Victoria, Australia), Hugh Mitenko (WA Country Health Service, Western Australia), Michael Lovegrove (Joondalup Health Campus, Western Australia), Ben Smedley (Rockingham General Hospital, Western Australia), Colin A Graham and Ling Yan Leung (Prince of Wales Hospital, Hong Kong SAR), Win Sen Kuan and Ying Wei Yau (National University Hospital, Singapore), Wei Ming Ng (Ng Teng

Fong General Hospital, Singapore), Ranjeev Kumar (Khoo Teck Puat Hospital, Singapore), Dennis Wen Jie Chia (Sengkang General Hospital, Singapore), Said Laribi (CHU Tours, Tours, France), Mounir Hilal and Rarthana Mil (CH Vendôme, France); Audrey Gerineau (CHR Orléans, France), Matthew J Reed (Emergency Medicine Research Group Edinburgh [EMERGE], Royal Infirmary of Edinburgh, United Kingdom), Daniel Horner (Salford Royal NHS Foundation Trust, Salford, United Kingdom), Edward Carlton and Tom Roberts (North Bristol NHS Trust, United Kingdom), Girish Boggaram and Jayne Foot (Musgrove Park Hospital, Taunton, United Kingdom), Andy Appleboam, Rachel Goss and Hamza Malik (Royal Devon and Exeter NHS Foundation, United Kingdom), Richard Body (Manchester Royal Infirmary, Manchester, United Kingdom), John-Paul Williamson (Royal Oldham Hospital, Oldham, United Kingdom); Adela Golea and Sonia Luka (University County Hospital Cluj-Napoca, Romania), Huseyin Avni Demir (University of Health Sciences Mehmet Akif Inan Training and Research Hospital, Department of Emergency Medicine, Şanlıurfa Turkey), Şafak Öner Gülpinar (Tokat Erbaa Government Hospital, Tokat, Turkey), Lale Tolu (Bursa Çekirge Government Hospital of Emergency Service, Bursa, Turkey), Muhammet Hacimustafaoğlu (Hakkari Yuksekova Government Hospital, Hakkari, Turkey), Mehmet A Karamercan (Gazi University Faculty of Medicine Department of Emergency Medicine, Ankara, Turkey), Elif Çelikel (Numune Research and Training Hospital, Department of Emergency Medicine, Ankara, Turkey), Çilem Çaltılı (University Of Health Sciences Bağcılar Training and Research Hospital, Department of Emergency Medicine, Istanbul, Turkey); Selahattin Gürü (Yıldırım Beyazıt University Faculty of Medicine Department of Emergency Medicine, Ankara, Turkey), Gülşah Yavuz (Antalya Ataturk Government Hospital of Medicine, Department of Emergency Medicine, Antalya, Turkey), Franck Verschuren (Institute of Experimental and Clinical Research, Emergency Department, Saint-Luc University Hospital, Brussels, Belgium), Christopher Ramos (Emergency Department, Saint-Luc University Hospital, Brussels, Belgium), Paule Denoel and Nicolas Wilmet (Saint Michel, Clinique de l'Europe, Etterbeek, Brussels), Michael Vandoorslaert and Alessandro Manara (Saint Elisabeth, Clinique de l'Europe, Uccle, Brussels), Adeline Higuët (CHR Hal, Belgium), Amichai Sheffy (Tel-Aviv Sourasky Medical Center, Israel), Sinan Kamona and Peter Jones (University of Auckland, School of Medicine, Auckland, New Zealand), Mai Nguyen (Wellington Hospital, Wellington, New Zealand); Anne Clarke (Hutt Valley Hospital, Lower Hutt, New Zealand), Sierra Beck (Dunedin Hospital, Dunedin, New Zealand), Andrew Munro (Nelson Hospital, Nelson, New Zealand), Kim M

Yates (North Shore and Waitakere Hospitals, Waitematā District Health Board, New Zealand), James Weaver (Christchurch Hospital, Christchurch, New Zealand), Deborah Moore and Stuart Innes (Tauranga Hospital, Tauranga, New Zealand), Karina Walters (Taranaki District Health Board, New Zealand), Koen Simons (Statistician, Office for Research, Western Health and University of Melbourne, Victoria, Australia); Steering Committee: Anne-Maree Kelly, Kevin Chu, Tissa Wijeratne, Frances B Kinnear, Gerben Keijzers, Sinan Kamona, Win Sen Kuan, Colin Graham, Richard Body, Said Laribi, Sharon Klim, Mehmet Karamercan, Tom Roberts; and co-ordinating Centre, Western Health, Victoria, Australia: Anne-Maree Kelly, Sharon Klim and Kerrie Russell.

### CONFLICTS OF INTEREST

No conflicts of interest to declare.

### REFERENCES

- Burch R, Rizzoli P, Loder E. The prevalence and impact of migraine and severe headache in the United States: Updated age, sex, and socioeconomic-specific estimates from government health surveys. *Headache*. 2021;61(1):60-68. doi:10.1111/head.14024
- Chu KH, Howell TE, Keijzers G, et al. Acute headache presentations to the emergency department: a statewide cross-sectional study. *Acad Emerg Med*. 2017;24(1):53-62. doi:10.1111/acem.13062.3
- Friedman BW, Serrano D, Reed M, et al. Use of the emergency department for severe headache. A population-based study. *Headache*. 2009;49(1):21-30. doi:10.1111/j.1526-4610.2008.01282.x
- Solomon GD, Kunkel RS Jr, Frame J. Demographics of headache in elderly patients. *Headache*. 1990;30(5):273-276. doi:10.1111/j.1526-4610.1990.hed3005273.x
- Ruiz M, Pedraza MI, de la Cruz C, et al. Headache in the elderly: characteristics in a series of 262 patients. *Neurologia*. 2014;29(6):321-326. doi:10.1016/j.nrl.2013.07.007
- Pascual J, Berciano J. Experience in the diagnosis of headaches that start in elderly people. *J Neurol Neurosurg Psychiatry*. 1994;57(10):1255-1257. doi:10.1136/jnnp.57.10.1255
- Wijeratne T, Tang HM, Crewther D, Crewther S. Prevalence of Migraine in the Elderly: A Narrated Review. *Neuroepidemiology*. 2019;52(1-2):104-110. doi:10.1159/000494758
- Do TP, Remmers A, Schytz HW, et al. Red and orange flags for secondary headaches in clinical practice: SNNOOP10 list. *Neurology*. 2019;92(3):134-144. doi:10.1212/WNL.00000000000006697
- Perry JJ, Stiell IG, Sivilotti ML, et al. Clinical decision rules to rule out subarachnoid hemorrhage for acute headache. *JAMA*. 2013;310(12):1248-1255. doi:10.1001/jama.2013.278018
- Ramirez-Lassepas M, Espinosa CE, Cicero JJ, Johnston KL, Cipolle RJ, Barber DL. Predictors of intracranial pathologic findings in patients who seek emergency care because of headache. *Arch Neurol*. 1997;54(12):1506-1509. doi:10.1001/archneur.1997.00550240058013
- Locker TE, Thompson C, Rylance J, Mason S. The utility of clinical features in patients presenting with nontraumatic headache: an investigation of adult patients attending an emergency department. *Headache*. 2006;46(6):954-961. doi:10.1111/j.1526-4610.2006.00448.x
- Munoz-Ceron J, Marin-Careaga V, Peña L, Mutis J, Ortiz G. Headache at the emergency room: Etiologies, diagnostic usefulness of the ICHD 3 criteria, red and green flags. *PLoS One*. 2019;14(1):e0208728. doi:10.1371/journal.pone.0208728
- Giamberardino MA, Affaitati G, Costantini R, Guglielmetti M, Martelletti P. Acute headache management in emergency department. *A narrative review. Intern Emerg Med*. 2020;15(1):109-117. doi:10.1007/s11739-019-02266-2
- Edlow JA, Panagos PD, Godwin SA, Thomas TL, Decker WW. American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with acute headache. *Ann Emerg Med*. 2008;52(4):407-436. doi:10.1016/j.annemergmed.2008.07.001
- Australasian College of Emergency Medicine. Guidelines on Diagnostic Imaging. <https://acem.org.au/getmedia/b7f67701-7e80-4e06-b87e-9193310415b0/Guidelines-for-Diagnostic-Imaging-Jul-12-v01.aspx>. Accessed 17.01.2021.
- Kelly AM, Kwan WS & Chu KH et al. Epidemiology, Investigation, Management and Outcome of Headache in Emergency Departments, (HEAD Study) – An Multinational Observational Cohort Study. *Headache* 2021 (at press)
- Orimo H, Ito H, Suzuki T, Araki A, Hosoi T, Sawabe M. Reviewing the definition of “elderly”. *Geriatr Gerontol Int*. 2006;6(3):149-158. doi:10.1111/j.1447-0594.2006.00341.x
- Ouchi Y, Rakugi H, Arai H, et al. Redefining the elderly as aged 75 years and older: Proposal from the Joint Committee of Japan Gerontological Society and the Japan Geriatrics Society. *Geriatr Gerontol Int*. 2017;17(7):1045-1047. doi:10.1111/ggi.13118
- von Elm E, Altman DG, Egger M, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007;335(7624):806-808. doi:10.1136/bmj.39335.541782.AD
- <https://www.stat.ubc.ca/~rollin/stats/ssize/b1.html>. Accessed July 2021
- Straube A, Andreou A. Primary headaches during lifespan. *J Headache Pain*. 2019;20(1):35. doi:10.1186/s10194-019-0985-0
- Aiello A, Farzaneh F, Candore G, et al. Immunosenescence and Its Hallmarks: How to oppose aging strategically? A review of potential options for therapeutic intervention. *Front Immunol*. 2019;10:2247. doi:10.3389/fimmu.2019.02247
- Canto JG, Shlipak MG, Rogers WJ, et al. Prevalence, clinical characteristics, and mortality among patients with myocardial infarction presenting without chest pain. *JAMA*. 2000;283(24):3223-3229. doi:10.1001/jama.283.24.3223
- Samaras N, Chevalley T, Samaras D, Gold G. Older patients in the emergency department: a review. *Ann Emerg Med*. 2010;56(3):261-269. doi:10.1016/j.annemergmed.2010.04.015
- Limpawattana P, Phungoen P, Mitsungnorn T, Laosuankoon W, Tansanhorn N. Atypical presentations of older adults at the emergency department and associated factors. *Arch Gerontol Geriatr*. 2016;62:97-102. doi:10.1016/j.archger.2015.08.016
- Kehoe A, Smith JE, Bouamra O, Edwards A, Yates D, Lecky F. Older patients with traumatic brain injury present with a higher GCS score than younger patients for a given severity of injury. *Emerg Med J*. 2016;33(6):381-385. doi:10.1136/emerm ed-2015-205180

27. Doretti A, Shestaritc I, Ungaro D, et al. Headaches in the emergency department -a survey of patients' characteristics, facts and needs. *J Headache Pain*. 2019;20(1):100. doi:10.1186/s10194-019-1053-5
28. Gilbert EH, Lowenstein SR, Koziol-McLain J, Barta DC, Steiner J. Chart reviews in emergency medicine research: Where are the methods? *Ann Emerg Med*. 1996;27(3):305-308. doi:10.1016/s0196-0644(96)70264-0

### SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Beck S, Kinnear FB, Maree Kelly A, et al; HEAD Study Group. Clinical presentation and assessment of older patients presenting with headache to emergency departments: A multicentre observational study. *Australas J Ageing*. 2021;00:1–12. <https://doi.org/10.1111/ajag.12999>